

STABILITY OF ITERATIVE PROCEDURES FOR HYBRID MAPS IN b-METRIC SPACE

AMAL M. HASHIM

Department of Mathematics, College of Science, University of Basrah, Basrah, Iraq

E-mail: amalmhashim@yahoo.com

Abstract.

In this paper, we study the existence of coincidences and fixed points of hybrid contractions maps on more general setting than metric spaces.

The same utilize to discuss the problem of stability of iterative procedures in generalized Hausdorff metric space.

Our work generalize and improve some results due to Ostrowski 1967, Czerwik et al. 1997, Singh et al 2005 and Singh et al. 2008.

Mathematics Subject Classifications (2000): 41A25, 65D15, 47H10

Key words and phrases: Stable iterative, coincidence point, hybrid maps, b-metric space.

1. INTRODUCTION

Let (X, d) be a metric space and an operator $T: X \rightarrow CL(X)$ where $CL(X)$ the collection of all nonempty closed subsets of X . Problem of existence of coincidence and fixed points of T lead to an iteration procedure $x_{n+1} \in f(T, x_n)$. However, in

computation, an approximate sequence $\{y_n\}$ is applied instead of $\{x_n\}$. This leads to a concept of stability of iteration procedure with respect to T .

The first stability result in metric space was proved by Ostrowski (1967). For

any $x_0 \in X$, we put
 $x_{n+1} = f(T, x_n), n = 1, 2, \dots$ (1)

Let $\{x_n\}$ be a sequence converges
 to a fixed point $u \in X$ of T . Let
 $\{y_n\} \subseteq X$ be an arbitrary sequence, and
 set

$$\varepsilon_n = d(y_{n+1}, Ty_n), \quad n = 0, 1, 2, \dots \quad (2).$$

The iteration procedure (1) is said to
 be T -stable (cf Czerwik 1998) provided
 that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ implies that $\lim_{n \rightarrow \infty} y_n = u$.

This subject was developed by
 Harder and Hicks 1988a, b, Rhoades 1993,
 Osilike 1995, Osilike 1996 and Berinde 2002
 (see also Ćirić 1972, Ćirić 1974, Czerwik
 1997, Czerwik et al. 2002, Singh & Chanda
 1995, Singh & Prasad 2008, Singh et al.
 2005 and Singh & Hashim 2005).

Czerwik 1997 and Czerwik et al.
 2002 has been extended Ostrowski's
 classical theorem for the stability of iterative
 procedures for multi-valued maps in b-
 metric spaces. The purpose of this paper is
 to study the stability of iteration process for
 hybrid contraction mappings involving
 single-valued and multi-valued maps in b-
 metric spaces.

2. PRELIMINARIES

Consistent with Czerwik 1998 and
 Singh & Hashim 2005, we use the
 following notations and definitions.

Definition 2.1: Czerwik 1998.

Let X be a nonempty set and $s \geq 1$ be a
 given real number. A function
 $d : X \times X \rightarrow R_+$ is a b-metric provided
 that for all $x, y, z \in X$,

$$d(x, y) = 0 \text{ iff } x = y, \quad (3)$$

$$d(x, y) = d(y, x), \quad (4)$$

$$d(x, z) \leq s[d(x, y) + d(y, z)] \quad (5)$$

The pair (X, d) is called a b-metric
 spaces.

Notice that the class of b-metric spaces is
 effectively larger than that of metric spaces.
 The following example shows that a b-
 metric on X need not a metric on X (see
 also Singh & Prasad 2008).

Example 2.1: Let $X = \{x_1, x_2, x_3\}$ and
 $d : X \times X \rightarrow R_+$ such that

$$d(x_1, x_2) = a \geq 2, \quad d(x_1, x_3) = d(x_2, x_3) = 1 \text{ and}$$

$$d(x_n, x_n) = 0, \quad d(x_n, x_k) = d(x_k, x_n),$$

$$d(x_n, x_k) = \frac{a}{2} [d(x_n, x_i) + d(x_i, x_k)], \quad n, k, i = 1, 2, 3$$

Then (X, d) is a b-metric space.

$CL(X) = \{A : A \text{ is nonempty closed}$
 subset of $X\}$

$$H(A, B) = \max \begin{cases} \sup_{x \in A} D(x, B), \sup_{y \in B} D(y, A), & \text{if the maximum exists} \\ \infty, & \text{otherwise} \end{cases}$$

$$A, B \in CL(X)$$

H is called a generalized Hausdorff b-metric on $CL(X)$

$$D(x, A) = \inf_{\alpha \in A} d(x, \alpha)$$

Lemma 2.1 Czerwik 1998:

For any $A, B, C \in CL(X)$,

$$D(x, B) \leq d(x, y) \text{ For any } y \in B, \quad (6)$$

$$D(A, B) \leq H(A, B), \quad (7)$$

$$H(A, C) \leq s[H(A, B) + H(B, C)]. \quad (8)$$

Lemma 2.2 (Czerwik 1998):

Let A and B be distinct elements of $CL(X)$

Let $x \in A$. Then for a given number $\lambda > 1$, there exists a point y in B such that $d(x, y) \leq \lambda H(A, B)$.

Definition 2.2 (Singh & Chanda 1995):

Let X be a metric space and $T : X \rightarrow CL(X)$. Let $x_0 \in X$, and

$$(*) x_{n+1} \in f(T, x_n),$$

denote some iteration procedure. Let the sequence $\{x_n\}$ be convergent to a fixed point P of T . Let $\{y_n\}$ be an arbitrary sequence in X and set

$$\varepsilon_n = H(y_{n+1}, f(T, y_n)), \quad n = 0, 1, 2, \dots,$$

If limit $\varepsilon_n = 0$ implies that $\lim_{n \rightarrow \infty} y_n = p$ then the iteration process defined in (*) is said to be T -stable or stable with respect to T .

Recall that this definition for a single-valued operator is due to Harder & Hicks 1988a, b.

The following definition of stability general iterative procedure (9) for the coincidence point equation $fx = Tx$ is due to Singh et al. 2008 when S and T are single-valued maps.

Definition 2.3 [Singh et al. 2008]:

Let (X, d) be a b-metric space and $Y \subseteq X$. Let $T : Y \rightarrow CL(X)$, $S : Y \rightarrow X$, $T(Y) \subseteq S(Y)$ and z a coincidence point of T and S , that is $Sz \in Tz = u$. For any $x_0 \in Y$, let the sequence $\{Sx_n\}$ generated by the general iterative procedure.

$$Sx_{n+1} \in f(T, x_n) \quad n = 1, 2, \dots \quad (9)$$

Converge to an element $u \in X$. Let $\{Sy_n\} \subseteq X$ be an arbitrary sequence and let $\varepsilon_n = H(Sy_{n+1}, f(T, y_n)), \quad n = 0, 1, 2, \dots$.

Then the iterative procedure $f(T, x_n)$ is (S, T) , stable or stable with respect to (S, T) if $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ implies that $\lim_{n \rightarrow \infty} S y_n = 0$.

We remark that the iterative procedure (9) reduce to $x_{n+1} \in f(T, x_n)$ $n = 1, 2, \dots$.

When $Y = X$ and S is the identity map on X . The general definition of stability of iterative procedures for general fixed equation due to Czerwik et al. 2002 and Singh et al. 2005.

Consider the following conditions for $T : Y \rightarrow CL(X)$ and $S : Y \rightarrow X$ for all $x, y \in Y$, where $0 < q < 1$ and X is a metric space;

$$H(Tx, Ty) \leq qd(Sx, Sy); \quad (10)$$

$$H(Tx, Ty) \leq q \max\{d(Sx, Sy), D(Sx, Tx), D(Sy, Ty), \frac{1}{2}[D(Sx, Ty) + d(Sy, Tx)]\}; \quad (11)$$

$$H(Tx, Ty) \leq q d(Sx, Sy) + LD(Sx, Tx). \quad 0 < q < 1, L \geq 0 \quad (12)$$

Notice that (10) is included in (11) while (11) itself is called a generalized hybrid contraction and when $S = id$ identity map on X (11) is called a generalized multi-valued contraction. These conditions with $X = Y$, $S = id$, the identity map on X and T is a single valued mapping are compared in Rhoades 1977.

Lemma 2.3: see also (Rhoades 1993, Czerwik et al. 2002).

Let $\{\varepsilon_n\}$ be a sequence of non negative real numbers.

$$\text{Let } s_n = \sum_{i=0}^n k^{n-i} \varepsilon_i \quad \text{where } 0 \leq k < 1.$$

$$\text{Then } \lim_{n \rightarrow \infty} \varepsilon_n = 0 \text{ iff } \lim_{n \rightarrow \infty} s_n = 0. \quad (13)$$

Now we can present the following fixed point theorem (see Singh et al. 2005)

Theorem 2.1: (Singh et al. 2005):

Let X be a complete b-metric space and T a generalized multi-valued contraction on X . Then

(i) For every $x_0 \in X$ there exists an orbit $\{x_n\}$ of T at x_0 and $u \in X$ such that $\lim_{n \rightarrow \infty} x_n = u$;

(ii) The point u is fixed point of T , i.e. $u \in Tu$.

3. Main Results

Theorems 3.1:

let (X, d) be a b-metric and $T : Y \rightarrow CL(X)$, $S : Y \rightarrow X$ such that $T(Y) \subseteq S(Y)$ and (11) holds with $qs < 1$ and $\alpha s < 1$ where

$$\alpha = \max\left\{k, \frac{ks}{2 - qs}\right\} < 1 \quad \text{and one of}$$

TY or SY is a complete subspace of X , then T and S have a coincidence point, i.e. there exists a $v \in Y$ $\ni Sv \in Tv$.

Proof:

Let $x_o \in Y$ since $T(Y) \subseteq S(Y)$, choose x_1 , so that $y_1 = Sx_1 \in Tx_o$. In general choose x_{n+1} so that $y_{n+1} = Sx_{n+1} \in Tx_n$. let λ be a real number such that $0 < \lambda < 1$. By lemma (2.2) and condition (11),

$$\begin{aligned} d(y_n, y_{n+1}) &\leq q^{-\lambda} H(Tx_{n-1}, Tx_n) \\ &\leq q^{-\lambda} \cdot q \max\{d(Sx_{n-1}, Sx_n), \\ &\quad D(Sx_{n-1}, Tx_{n-1}), D(Sx_n, Tx_n), \\ &\quad \frac{1}{2}[D(Sx_{n-1}, Tx_n) + D(Sx_n, Tx_{n-1})]\} \\ d(y_n, y_{n+1}) &\leq k \max\{d(y_{n-1}, y_n), d(y_{n-1}, y_n), \\ &\quad d(y_n, y_{n+1}), \frac{1}{2}[d(y_{n-1}, y_{n+1}) + d(y_n, y_n)]\} \end{aligned}$$

where $k = q^{1-\lambda}$.

Since $k < 1$ the relation

$$d(y_{n-1}, y_{n+1}) \leq kd(y_n, y_{n+1}) \text{ implies}$$

$$d(y_{n-1}, y_{n+1}) = 0.$$

We may assume that $y_n \neq y_{n+1}$ and

$$d(y_n, y_{n+1}) \leq k \max\{d(y_{n-1}, y_n), \frac{1}{2}d(y_{n-1}, y_{n+1})\}$$

so $d(y_n, y_{n+1}) \leq kd(y_{n-1}, y_n)$, or else

$$d(y_n, y_{n+1}) \leq \frac{ks}{2} [d(y_{n-1}, y_n) + d(y_n, y_{n+1})],$$

$$\text{so } d(y_n, y_{n+1}) \leq \frac{ks}{2-ks} d(y_{n-1}, y_n)$$

This yield

$$d(y_n, y_{n+1}) \leq \max\{k, \frac{ks}{2-ks}\} [d(y_{n-1}, y_n)] \quad n=1, 2, \dots,$$

$$\text{where } \max\{k, \frac{ks}{2-ks}\} < 1.$$

Therefore, $\{y_n\}$ is a Cauchy sequence and its sub sequence $\{y_{2n}\}$ has a limit in SY

call it u . let $v \in S^{-1}u$. Then $Sv = u$.

Notice that the subsequence $\{y_{2n+1}\}$ also converge to u . by (11)

$$\begin{aligned} D(Tv, y_{2n+2}) &\leq H(Tv, Tx_{2n+1}) \\ &\leq q \max\{d(Sv, Sx_{2n+1}), D(Sv, Tv), D(Sx_{2n+1}, Tx_{2n+1}) \\ &\quad \frac{1}{2}[D(Sv, Tx_{2n+1}) + D(Sx_{2n+1}, Tv)]\}, \\ &\leq q \max\{d(Sv, y_{2n+1}), D(Sv, Tv), d(y_{2n+1}, y_{2n+2}), \\ &\quad \frac{1}{2}[d(Sv, y_{2n}) + D(y_{2n+1}, Tv)]\}. \end{aligned}$$

Making $n \rightarrow \infty$ this obtains

$$D(Tv, Sv) \leq q D(Sv, Tv), \text{ and } Sv \in Tv$$

We remark that theorem (3.1) with $S = Id$ (the identity map) is the main result of Singh et al. 2005.

If T is a single-valued map in theorem 3.1 with $s=1$ is also the main result of Ciric 1972.

Theorem 3.2: let (X, d) be a b-metric and $Y \subseteq X$ let $T: Y \rightarrow CL(X)$, $S: Y \rightarrow X$ and SY or TY is a complete subspace of X .

Let z be a coincidence point of T and S , i.e., $u = Sz \in Tz$. For any $x_0 \in Y$, let the sequence $\{Sx_n\}$, generated by $Sx_{n+1} \in Tx_n$ converge to u . Let $\{Sy_n\} \subseteq X$ and define $\varepsilon_n = H(Sy_{n+1}, Ty_n)$, $n = 0, 1, 2, \dots$. If S and T satisfy (11) for all $x, y \in Y$ and $k = sq < 1$, then

$$d(u, Sy_{n+1}) \leq sd(u, Sx_{n+1}) + s(s\alpha)^{n+1}d(Sx_0, Sy_0) + s^3\alpha \sum_{r=0}^n (s\alpha)^{n-r}d(Sx_n, Sx_{n+1}) + s^2 \sum_{r=0}^n (s\alpha)^{n-r}\varepsilon_r \quad (14)$$

$$\alpha = s^2q / 1 - s^2q$$

$$\lim_{n \rightarrow \infty} Sy_n = u \Leftrightarrow \lim_{n \rightarrow \infty} \varepsilon_n = 0 \quad (15)$$

Proof: From (11) for any $x, y \in Y$, one of the following holds;

$$\begin{aligned} H(Tx, Ty) &\leq qd(Sx, Sy), \\ H(Tx, Ty) &\leq qD(Sx, Tx) \leq qH(Sx, Tx); \\ H(Tx, Ty) &\leq qD(Sy, Ty) \leq qH(Sy, Ty) \leq qs[H(Sy, Sx) + H(Sx, Ty)] \\ &\leq qsD(Sy, Sx) + qs[sH(Sx, Tx) + sH(Tx, Ty)] \\ &\leq qsD(Sy, Sx) + qs^2H(Sx, Tx) + qs^2H(Tx, Ty) \end{aligned}$$

$$\text{i.e., } H(Tx, Ty) \leq \frac{qs}{1-qs^2}d(Sy, Sx) + \frac{qs^2}{1-qs^2}H(Sx, Tx)$$

$$\leq \frac{k}{1-ks}d(Sy, Sx) + \frac{ks}{1-ks}H(Sx, Tx)$$

$$\begin{aligned} H(Tx, Ty) &\leq \frac{q}{2}\{D(Sx, Ty) + D(Sy, Tx)\} \\ &\leq \frac{q}{2}\{H(Sx, Ty) + H(Sy, Tx)\} \\ &\leq \frac{qs}{2}\{H(Sx, Tx) + H(Tx, Ty) + H(Sy, Sx) + H(Sx, Tx)\} \\ &\leq \frac{qs}{2}\{2H(Sx, Tx) + H(Sx, Sy) + H(Tx, Ty)\} \end{aligned}$$

i.e.,

$$\begin{aligned} H(Tx, Ty) &\leq \frac{qs}{2-qs}\{2H(Sx, Tx) + d(Sx, Sy)\} \\ &\leq \frac{k}{2-k}d(Sx, Sy) + \frac{2k}{2-k}H(Sx, Tx) \end{aligned}$$

Therefore, in all the cases we get,

$$H(Tx, Ty) \leq \alpha d(Sx, Sy) + s\alpha H(Sx, Tx)$$

$$\text{Where } \alpha = \frac{s^2q}{1-s^2q}$$

Then by virtue of (11) we obtain, For any non-negative integer n .

$$\begin{aligned} d(Sx_{n+1}, Sy_{n+1}) &\leq H(Tx_n, Sy_{n+1}) \leq s[H(Tx_n, Ty_n) + H(Ty_n, Sy_{n+1})] \\ &\leq s[\alpha d(Sx_n, Sy_n) + s\alpha H(Sx_n, Tx_n) + \varepsilon_n] \\ &\leq s\alpha d(Sx_n, Sy_n) + s^2\alpha H(Sx_n, Tx_n) + s\varepsilon_n \\ &\leq s^2\alpha[H(Tx_{n-1}, Ty_{n-1}) + H(Ty_{n-1}, Sy_n)] + s^2\alpha H(Sx_n, Tx_n) + s\varepsilon_n \\ &\leq s^2\alpha[\alpha d(Sx_{n-1}, Sy_{n-1}) + \alpha sd(Sx_{n-1}, Tx_{n-1})] + \\ &\quad s^2\alpha\varepsilon_{n-1} + s^2\alpha H(Sx_n, Tx_n) + s\varepsilon_n \\ &\leq s^2\alpha^2 d(Sx_{n-1}, Sy_{n-1}) + (s^2\alpha^2)sd(Sx_{n-1}, Tx_{n-1}) + \\ &\quad s^2\alpha\varepsilon_{n-1} + s\varepsilon_n + s^2\alpha H(Sx_n, Tx_n) \\ &\leq (s\alpha)^2 d(Sx_{n-1}, Sy_{n-1}) + (s\alpha)^2 sd(Sx_{n-1}, Tx_{n-1}) + \\ &\quad s^2\alpha H(Sx_n, Tx_n) + s[\alpha s\varepsilon_{n-1} + \varepsilon_n] \end{aligned}$$

Repeating this process $(n-1)$ times we get

$$d(Sx_{n+1}, Sy_{n+1}) \leq (s\alpha)^{n-1}d(Sx_0, Sy_0) + s^2\alpha \sum_{r=0}^n (s\alpha)^{n-r}d(Sx_r, Ty_r) + s \sum_{r=0}^n (s\alpha)^{n-r}\varepsilon_r$$

Consequently,

$$d(u, Sy_{n+1}) \leq sd(u, Sx_{n+1}) + sd(Sx_{n+1}, Sy_{n+1})$$

This yields the inequality (14).

Assume that $\lim_{n \rightarrow \infty} Sy_n = u$

$$\begin{aligned} \varepsilon_n = H(Sy_{n+1}, Ty_n) &\leq sH(Sy_{n+1}, u) + sH(u, Ty_n) \\ &\leq sd(Sy_{n+1}, u) + s[\alpha d(u, Sy_n) + s\alpha H(Su, Tu)] \end{aligned}$$

Letting $n \rightarrow \infty$ we obtain $\varepsilon_n \rightarrow 0$

Suppose $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ since $0 \leq s\alpha < 1$ and

$$\lim_{n \rightarrow \infty} Sx_n = u$$

First two terms on the right hand side (14) vanish in the limit. Applying lemma (2.3) to the last term of inequality (14) we see that

$$\lim_{n \rightarrow \infty} s^2 \sum_{r=0}^{\infty} (s\alpha)^{n-r} \varepsilon_r = 0$$

Finally we show that,

$$\lim_{n \rightarrow \infty} s^3 \alpha \sum_{r=0}^n (s\alpha)^{n-r} d(Sx_r, Tx_r) = 0$$

Let A denote the lower triangular matrix with entries $a_{nr} = (s\alpha)^{n-r}$, then

$$\lim_{n \rightarrow \infty} a_{nr} = 0 \quad \text{for each } r \quad \text{and}$$

$$\lim_{n \rightarrow \infty} \sum_{r=0}^n a_{nr} = \frac{1}{1-s\alpha} \quad \text{therefore, } A \text{ is}$$

multiplicative i.e for any converge sequence

$$\{s_n\}, \lim_{n \rightarrow \infty} A(s_n) = \frac{1}{1-s\alpha} \lim_{n \rightarrow \infty} s_n. \text{ thus,}$$

$$\lim_{n \rightarrow \infty} s^3 \alpha \sum_{r=0}^n (s\alpha)^{n-r} d(Sx_r, Tx_r) = 0$$

(cf. Czerwik 1997 p. 692).

Remark 3.1:

Theorem (3.2) with $S = I$ (identity map) generalize the main result of Singh et al. 2005.

Corollary 3.1: Singh et al. 2005 .

Let (X, d) be a complete b-metric space and $T : X \rightarrow CL(X)$ satisfy condition (11) with $S = I$ (identity map), let $x_o \in X$ and $\{x_n\}$ be an orbit for T at

x_n , i.e. $x_{n+1} \in Tx_n \quad n = 0, 1, 2, \dots$

and $\{x_n\}$ converge to a fixed point u of

T . Let $\{y_n\}$ be a sequence in X and set

$$\varepsilon_n = H(y_{n+1}, Ty_n), \quad n = 0, 1, 2, \dots$$

then

$$d(u, Sy_{n+1}) \leq s(sq)^{n+1} d(x_o, y_o) + s^2 m q \sum_{r=0}^n (sq)^{n-r} d(x_r, x_{r+1}) + s^2 \sum_{r=0}^n (sq)^{n-r} \varepsilon_r$$

Where $m = (1 - sq)^{-1}$

If Tu is singleton, then $\lim_{n \rightarrow \infty} y_n = u$ iff

$$\lim_{n \rightarrow \infty} \varepsilon_n = 0$$

Corollary 3.2: (Czerwik et al. 2002)

Let (X, d) be a complete b-metric space and let $T : X \rightarrow CL(X)$ satisfy

$$H(Tx, Ty) \leq \alpha d(x, y), \quad x, y \in \lambda$$

where $0 < \alpha < s^{-1}$ let $x_o \in X$ and $\{x_n\}$

be an orbit of T at x_o . i.e.

$$x_{n+1} \in Tx_n, \quad n = 0, 1, 2, \dots \text{ and } \{x_n\}$$

converge to fixed point u of T , moreover

let $\{y_n\}$ be a sequence in X and set

$$\varepsilon_n = H(y_{n+1}, Ty_n), \quad n = 0, 1, 2, \dots$$

then

$$d(u, y_{n+1}) \leq sd(u, x_{n+1}) + s(s\alpha)^{n+1} d(x_o, y_o) + s^2 \sum_{r=0}^n (s\alpha)^{n-r} \varepsilon_r$$

if moreover, Tu is singleton, then

$$\lim_{n \rightarrow \infty} y_n = u \text{ iff } \lim_{n \rightarrow \infty} \varepsilon_n = 0$$

Remark 3.2: 1- Theorem (3.2) generalize theorem 4.3 of Singh et al. 2008. when S and T are single valued maps .

2- If $s = 1$, $S = Id$ (the identity map), S and T are single valued map in the theorem 3.2, we obtain the main result of Osilike 1995 .

Theorem 3.3: let (X, d) be a b-metric and $Y \subseteq X$ let $T : Y \rightarrow CL(X)$, $S : Y \rightarrow X$ such that $T(Y) \subseteq S(Y)$ and $S(Y)$ or $T(Y)$ is a complete subspace of X . Let z be a coincidence point of T and S , i.e. $u = Sz \in Tz$ generated by $Sx_{n+1} \in Tx_n$ converge to u . Let $\{Sy_n\} \subset X$ and define

$$\varepsilon_n = H(Sy_n, Ty_n), n = 0, 1, 2, \dots$$

If S and T satisfies (12) with $s^2q < 1$, then

$$d(u, Sy_{n+1}) \leq sd(u, Sx_{n+1}) + s(sq)^{n+1}d(Sx_0, Sy_0) + \quad (16)$$

$$s^2L \sum_{r=0}^n (sq)^{n-r} d(Sx_r, Tx_r) + s^2 \sum_{r=0}^n (sq)^{n-r} \varepsilon_r.$$

$$\lim_{n \rightarrow \infty} Sy_n = u \Leftrightarrow \lim_{n \rightarrow \infty} \varepsilon_n = 0. \quad (17)$$

Proof : In view of the fact that $Sx_{n+1} \in Tx_n$ for $n = 0, 1, 2, \dots$ and the definition of H , we get $d(Sx_{n+1}, Sy_{n+1}) \leq H(Tx_n, Sy_{n+1})$.

Now, since H is a b-metric, so for any $A, B, C \in CL(X)$ we have

$$H(A, C) \leq s[H(A, B) + H(B, C)]$$

therefore by (12) we obtain

$$\begin{aligned} d(Sx_{n+1}, Sy_{n+1}) &\leq H(Tx_n, Sy_{n+1}) \leq s[H(Tx_n, Ty_n) + H(Ty_n, Sy_{n+1})] \\ &\leq s[qd(Sx_n, Sy_n) + LD(Sx_n, Tx_n)] + s\varepsilon_n \\ &\leq (sq)^2 d(Sx_{n-1}, Sy_{n-1}) + s^2 q L d(Sx_{n-1}, Tx_{n-1}) + \\ &\quad sL d(Sx_n, Tx_n) + s^2 q \varepsilon_{n-1} + s\varepsilon_n \end{aligned}$$

Since

Repeating this process $n - 1$ times, we get

$$\begin{aligned} d(u, Sy_{n+1}) &\leq sd(u, Sx_{n+1}) + s(sq)^{n+1} d(Sx_0, Sy_0) + s^2 L \sum_{r=0}^n (sq)^{n-r} d(Sx_r, Tx_r) + \\ &\quad s^2 \sum_{r=0}^n (sq)^{n-r} \varepsilon_r \end{aligned}$$

The proof (17) is similar to (15) in Theorem 3.2.

Remark 3.3:

1: In Theorem 3.3 when we put S and T maps on arbitrary set Y with values in X we get Theorem 4.2 in Singh and Prasad 2008. and when s generalize the main result of Singh et al. 2005.

2: Theorem 3.3 generalize the main result of Rhoades 1993 in case of single value map and when $s = 1$ and $S = Id$ (Identity map) under the following condition ;

$$d(Tx, Ty) \leq q \max\{d(x, y), \frac{d(x, Tx) + d(y, Ty)}{2}, d(x, Ty), d(y, Tx)\}$$

We remark that the above condition implies (12). For more detail see Rhoades 1977 .

REFERENCES

Berinde, Vasile, Iterative approximation of fixed points, Efemeride Publishing House, Romania, 2002.

1. Ciric Lj .B.,Fixed points for generalized multi-valued contractions ,Mat. Vesnik 9(24) (1972) , 265-272.
2. Ciric Lj .B.,,A generalization of Banach's contraction principle , Proc. Amer. Math. Soc. 45 (1974) 267-273.
3. Czerwik, Stefan, Nonlinear set-valued contraction mappings in b-metric spaces, Attisem.Mat. Fis. Univ .Modena 45(2) (1998) 263-276.
4. Czerwik, Stefan, Krzysztof Dłutek & S.L Singh, Round-off stability of iteration procedures for set -valued operators in b- metric spaces, J. Natur. Phys.Sci.15(2002) 1-8.
5. Czerwik, Stefan, Krzysztof Dłutek & S.L Singh, Round-off stability of iteration procedures for operators in b- metric spaces, J. Natur. Phys.Sci.11(1997) 87-94.
6. Harder A.M. & T.L. Hicks (a), A stable iteration procedure for non-expansive mappings. Math. Japon.33 (1988) 687-692.
7. Harder A.M. & T.L. Hicks (b), Stability results for fixed point iteration procedures, Math, Japon.33 (1988) 693-706.
8. Osilike, M.O. A stable iteration procedure for quasi- contractive maps, Indian J. Pure. Appl. Math.27(1) (1996) 25-34.
9. Osilike, M.O, Stability results of fixed point iteration procedures, J. Nigerian Math .Soc.14(1995) 17-2.
10. Ostrowski A.M., The round-off stability of iteration ,z Angew .Math. Mech. 47 (1967) 77-81.
11. Rhoades B.E, Fixed point theorems and stability results for fixed point iteration procedures ,Indian J. Pure. Appl. Math. 21 (1990) 1-.9.
12. Rhoades B.E, Fixed point theorems and stability results for fixed point iteration procedures ,Indian J. Pure. Appl. Math. .24(1993) 697-703.
13. Rhoades B.E, A comparison of various definitions of contractive mappings, Trans. Amer. Math. Soc.226 (1977) (257-290).
14. Singh S.L and Chadha, Round-off stability of iteration for multivalued operators, C.R. Math. Rep. Acad. Sci. Canada 17 (5) (1995) ,187-192.
15. Singh S.L, & Bhagwati Prasad, Some coincidence theorem and stability of iterative procedures, Computers and

- Mathematics with Applications 55 (2008) 2512-2520.
16. Singh S.L, Charu Bhatnagar & ,Amal .M.Hashim, Round-off stability of Picard iteration for multivalued operators ,Nonlinear Anal. Forum 10 (2005) 15-19.
17. Singh S.L, Charu Bhatnagar & S.N. Mishra, Stability of iterative Procedures for multivalued maps in b-metric spaces, Denonstratio. Math.38 (4) (2005) 905-916.
18. Singh S.L,Charu Bhatnagar & S.N Mishra, Stability of Jungck-type iterative procedures, Int. J. Math. Sci 19 (2005),3035-3043.
19. Singh S.L. & Amal M. Hashim, New coincidence and fixed point theorems for strictly contractive hybrid maps. Australian Journal of Mathematical Analysis and Applications. (2005) 1-7.

استقرار العمليات المكررة للدوال الهجينية في الفضاء المترى – ب

د. أمل محمد هاشم البطاط

قسم الرياضيات – كلية العلوم- جامعة البصرة – البصرة – العراق

الملخص

تضمن هذا البحث دراسة وجود النقاط الثابتة والمتطابقة للدوال الهجينية على فضاء أكثر تعميماً من الفضاء المترى وكذلك تم استخدام نفس الطريقة لمناقشة استقرار العمليات المكررة في الفضاء الهاوزدورفي المترى المعمم. في هذا البحث تم تعميم وتطوير نتائج العديد من الباحثين مثل, Ostrowski 1967, Czerwik et al. 1997, Singh et al. 2005 و Singh et al. 2008.
